POSITION PAPER



Leveraging causality and explainability in digital agriculture

Ilias Tsoumas^{1,2}, Vasileios Sitokonstantinou³, Georgios Giannarakis¹, Evagelia Lampiri⁴, Christos Athanassiou⁴, Gustau Camps-Valls³, Charalampos Kontoes¹ and Ioannis N. Athanasiadis²

¹BEYOND Centre, IAASARS, National Observatory of Athens, Athens, Greece

²Artificial Intelligence, Wageningen University & Research, Wageningen, The Netherlands

³Image Processing Laboratory (IPL), Universitat de València, València, Spain

⁴Department of Agriculture, Crop Production and Rural Environment, University of Thessaly, Volos, Greece

Corresponding authors: Ilias Tsoumas and Vasileios Sitokonstantinou; Emails: i.tsoumas@noa.gr;

vasileios.sitokonstantinou@uv.es

Received: 08 November 2024; Revised: 15 January 2025; Accepted: 20 March 2025

Keywords: agriculture; causal inference; evidence-based decision making; pest management; causal machine learning

Abstract

Sustainable agricultural practices have become increasingly important due to growing environmental concerns and the urgent need to mitigate the climate crisis. Digital agriculture, through advanced data analysis frameworks, holds promise for promoting these practices. Pesticides are a common tool in agricultural pest control, which are key in ensuring food security but also significantly contribute to the climate crisis. To combat this, Integrated Pest Management (IPM) stands as a climate-smart alternative. We propose a causal and explainable framework for enhancing digital agriculture, using pest management and its sustainable alternative, IPM, as a key example to highlight the contributions of causality and explainability. Despite its potential, IPM faces low adoption rates due to farmers' skepticism about its effectiveness. To address this challenge, we introduce an advanced data analysis framework tailored to enhance IPM adoption. Our framework provides (i) robust pest population predictions across diverse environments with invariant and causal learning, (ii) explainable pest presence predictions using transparent models, (iii) actionable advice through counterfactual explanations for in-season IPM interventions, (iv) fieldspecific treatment effect estimations, and (v) assessments of the effectiveness of our advice using causal inference. By incorporating these features, our study illustrates the potential of causality and explainability concepts to enhance digital agriculture regarding promoting climate-smart and sustainable agricultural practices, focusing on the specific case of pest management. In this case, our framework aims to alleviate skepticism and encourage wider adoption of IPM practices among policymakers, agricultural consultants, and farmers.

Impact Statement

We present a new data analysis framework based on causality and explainability to help farmers adopt sustainable alternatives to traditional practices for agricultural management. The framework makes agricultural management more practical and trustworthy by providing clear, reliable predictions, advice tailored to specific fields, and impact assessment of recommended actions. In our example, this could lead to less reliance on harmful pesticides, helping to protect the environment and fight climate change. With this tool, farmers can make better-informed decisions that benefit their crops and the planet, promoting a healthier and more sustainable future.

© The Author(s), 2025. Published by Cambridge University Press. This is an Open Access article, distributed under the terms of the Creative Commons Attribution licence (http://creativecommons.org/licenses/by/4.0), which permits unrestricted re-use, distribution and reproduction, provided the original article is properly cited.

1. Introduction

Digital agriculture integrates agricultural expertise with digital technologies, such as remote sensing, IoT, and data analytics, to effectively leverage diverse data sources like satellite imagery, weather forecasts, and soil health metrics. This approach promotes more sustainable, resilient, and profitable farming by enabling data-driven decisions across the agricultural value chain (Basso and Antle, 2020). This approach is essential for adapting agriculture to our rapidly changing climate and mitigating its impact on climate change (Balasundram et al., 2023). Artificial Intelligence (AI) serves digital agriculture as the means to transform the data into insights, estimations, forecasts, and recommendations that aim to support decisionmaking to balance agriculture's environmental, societal, and economic aspects. However, digital agriculture has remained largely confined to using almost solely correlation-based AI, which excels at predictive tasks but cannot go further. In this context, we propose exploiting two underutilized branches of AI by digital agriculture-causality and explainability. They can unlock capabilities beyond the continuous pursuit of prediction accuracy for enhancing digital agriculture, given that it considers agricultural knowledge and practice and integrates it into the modeling and inference parts (Sitokonstantinou et al., 2024). Thus, causality and explainability bring in digital agriculture domainaware robust models, explainable predictions, counterfactual reasoning, and quantifying effects of advice, action, and policy.

Pest management is a quintessential example in this context, demonstrating the valuable contributions that causality and explainability offer. Conventional pest management has been shown to contribute to climate change. Raising temperatures, intensifying ultraviolet radiation, and reducing relative humidity are expected to increase pest outbreaks and undermine the efficacy of pest control methods like host-plant resistance, bio-pesticides, and synthetic pesticides (Sharma and Prabhakar, 2014; Skendžić et al., 2021). Despite climate experts' warnings, pesticide use in agriculture adversely affects public health (Boedeker et al., 2020) and contributes to the climate crisis. This impact includes: (i) greenhouse gas (GHG) emissions from pesticide production, packaging, and transportation (Audsley et al., 2009), (ii) compromised soil carbon sequestration (Xu et al., 2020), (iii) elevated GHG emissions from soil (Spokas and Wang, 2003; Marty et al., 2010; Heimpel et al., 2013), and (iv) contamination of adjacent soil and water ecosystems, resulting in biodiversity loss (Sharma et al., 2019).

Thus, a vicious cycle has been established between pesticides and climate change (Sharma et al., 2022). In response, the European Commission (EC) has taken action to reduce all chemical and highrisk pesticides by 50% by 2030. Achieving such reductions requires adopting integrated pest management (IPM), which promotes sustainable agriculture and agroecology. IPM consists of eight principles inspired by the Food and Agriculture Organization (FAO) description. The authors in Barzman et al. (2015) condense these principles into *prevention and suppression, monitoring*, *decision-making, non-chemical methods, pesticide selection, reduced pesticide use, anti-resistance strategies*, and *evaluation*.

Data-driven methods have played a crucial role in optimizing pest management decisions. Some studies employ supervised machine learning techniques, such as Random Forests and Artificial Neural Networks (ANNs), satellite Earth observations, and in-situ data for pest presence prediction (Aparecido et al., 2019; Zhang et al., 2019). Others extend their models to include weather data (Skawsang et al., 2019). Recurrent Neural Networks (RNNs) capture temporal features from weather data, effectively handling unobservable counterfactual outcomes (Xiao et al., 2019). Iost Filho et al. (2022) highlight the extraction of fine-scale information for Integrated Pest Management (IPM) using meteorological data, insect scouting records, machine learning, and remote sensing. Nanushi et al. (2022) propose an interpretable machine learning solution integrating numerical weather predictions, vegetation indices, and trap catch data for estimating *Helicoverpa armigera* presence in cotton fields. This approach enhances the decision-making aspect of IPM, shifting away from traditional threshold-based pesticide applications. The interpretability of these predictions enhances trust and allows for incorporating domain expertise in pest management decision-making.



Figure 1. Causal and explainable data analysis framework for enhanced IPM.

2. Proposal

As Barzman et al. (2015) point out, threshold-based and "spray/don't spray" advice is not enough. There is a need for a new class of digital tools that consider the entire set of IPM principles to enhance decision-making truly. In this direction, we propose a data analysis framework for IPM based on causality and explainability. It consists of short-term actionable advice for in-season interventions and long-term advice for supporting strategic farm planning (Figure 1).

This way, we will upgrade the *monitoring* and *decision-making* IPM principles leading to actionable advice for direct pest control interventions and assisting the selection of practices relevant to other IPM principles, such as the *use of non-chemical methods* and *reduce pesticide dosage*. Additionally, the proposed framework will better inform farmers concerning the potential impact of practices that, in turn, will enhance the IPM principle of *prevention and suppression*, for example, crop rotation, day of sowing, and no-tillage. Furthermore, our framework employs observational causal inference to continuously assess the recommendations above and satisfy the IPM principle of *evaluation*.

In this study, we exploit the proposed framework, demonstrating its applicability and efficiency in a case study for pest management. While the case study is specific it represents the general case of pest management in several crops and conditions, and the typical availability of data for such case studies.

3. Data

Our approach relies on diverse data sources as a key leverage to capture a comprehensive picture of the past, present, and future agro-environmental conditions. This will enable us to improve the modeling and comprehension of pest dynamics.

3.1. Earth observations

We leverage biophysical and biochemical properties such as Leaf Area Index (LAI), Normalized Difference Vegetation Index (NDVI), chlorophyll content, as well as data on evapotranspiration and soil moisture. These factors play a crucial role in monitoring pest population dynamics. The data is derived from the Sentinel-1/2 and Terra/Aqua (MODIS) satellite missions that provide open access to optical multi-spectral and Synthetic Aperture Radar (SAR) images.

3.2. Terrain & soil characteristics

We incorporate data from open-access digital elevation models and information on topsoil physical properties and soil organic carbon content (de Brogniez et al., 2015; Ballabio et al., 2016). This allows us to include fixed or long-term characteristics specific to the area of interest.



Figure 2. Traps distribution in the Greek mainland for 2019–2022. Colors indicate the different agroclimatic zones in which traps from the dataset belong. These zones have been identified based on the study conducted by Ceglar et al. (2019).

3.3. Numerical weather predictions (NWP) and reanalysis of environmental datasets

Any high spatial resolution weather forecast can be used. We utilize a custom configuration of WRF-ARW (Skamarock et al., 2019) at a spatial resolution of 2 km. Hourly predictions are made, and for each trap location (i.e., where we have measurements about pest abundance), we obtain daily values for air (2 m) and soil temperature (0 m), relative humidity (RH), accumulated precipitation (AP), dew point (DP), and wind speed (WS). These parameters have been widely used in related work and are extremely valuable for learning from past (reanalysis) and future (NWP) pest states.

3.4. In-field measurements

In-field measurements involve ground observations of pest abundance using pheromone traps specifically designed for monitoring the cotton bollworm, known by the scientific name *Helicoverpa armigera* (*H. armigera*). These traps contain the active ingredients Z-11-hexadecen-1-al and Z-9-hexadecenal. The traps are used from the beginning of the first generation until the end of the season, with regular replacement every 4 to 6 weeks. The company Corteva Agriscience Hellas has established a dense (in time and space) trap network (Figure 2) that covers almost all areas in the Greek mainland where cotton is cultivated. The traps are strategically positioned at suitable distances from each other to prevent interference and ensure accurate data collection. An agronomist examines the traps and counts the trapped insects at regular intervals every 3–5 days. Corteva Agriscience Hellas provides historical data consisting of 398 trap sequences and 8202 unique data points from 2019 to 2022 (Table 1). They also provide auxiliary data on pesticide application, potential crop damage from pests, the severity of the damage, trap replacements, and scouter comments.

4. Approach and methods

4.1. Causal graph for representing domain knowledge

We constructed a causal graph (Figure 3) based on domain knowledge and expertise, denoted as G, that represents the underlying causal relationships within the pest-farm ecosystem for the *H. armigera* case.

Year	Traps	Measurements	Mean	std	Sprays	Sprayed fields %
2022	126	2507	19.73	4.22	30	18.25
2021	109	2245	20.30	1.79	17	11.01
2020	81	1693	20.54	4.77	12	8.64
2019	82	1757	21.29	6.43	21	21.95

 Table 1. Summary of trap data



Figure 3. Causal graph of a pest-farm ecosystem for Helicoverpa armigera case.

The graph G comprises vertices V, which represent the variables in the system, and directed edges E, which symbolize the cause-and-effect relationships between these variables. Besides helping us articulate domain knowledge, the causal graph G will benefit the downstream technical analyses in various ways. For instance, G will be employed for effect identification via graphical tests (Pearl, 2009), where the structure of G is integral to discerning causal relationships. Conversely, in the case of estimating conditional average treatment effects within the potential outcomes framework, G will be utilized as a conceptual guide for considering causal structures during the control phase. In invariant causal prediction, the graph will facilitate the construction of an accurate list of invariant features using causal parents of the target outcome. Moreover, the structural knowledge captured in G could benefit invariant learning methods by guiding the environment E definition. This diverse and tailored incorporation of G is aimed at optimizing the utilization of domain knowledge by the specifications and objectives of each analytical technique.

Specifically, in the current case of the pest-farm ecosystem of *H. armigera*, various biotic and abiotic factors (Table 2) can influence the population dynamics *Y* of *H. armigera* (Sharma et al., 2012). Temperature *T* plays a crucial role, affecting the insect's growth, development, fecundity, and survival (Howe, 1967). The size *SG* of the first generation is related to the size of the second generation, and the Southern Oscillation Index *SOI* has a significant correlation with the size of the first spring generation (Maelzer and Zalucki, 1999, 2000). Additionally, the life cycle *LC* of *H. armigera* is temperature-dependent, with completion occurring between 17.5°C and 32.5°C (Mironidis and Savopoulou-Soultani, 2014). The presence of parasitoids and natural enemies in cotton cultivation is crucial to many IPM programs, including the control of *H. armigera* (Pereira et al., 2019). Many egg parasitoids of different families are known for their high parasitism *P* rates and their effectiveness in reducing the population of *H. armigera* (Noor-ul-Ane et al., 2015). Nevertheless, parasitism rates are influenced by temperature and

Id	Variable description	
T	Temperature	
SW	Soil water	
RHa	Air relative humidity	
SG	Size of generation	
Pr	Precipitation	
LC	Life cycle	
Р	Parasitism	
V	Variety	
Sp	Spraying	
CS	Cropping system	
AC	Adjacent crops	
W	Wind	
Ws	Spraying wind	
SOI	South oscillation index	
PGS	Plant growth stage	
Y	Outcome (H. armigera population)	

 Table 2. Pest-farm ecosystem variables

relative humidity (Kalyebi et al., 2005; Noor-ul-Ane et al., 2015). Moreover, the efficacy of spray application Sp also impacts population dynamics (Wardhaugh et al., 1980). The efficacy of Sp is significantly influenced by the plant growth stage PGS. During the seedling stage, limited leaf surface area reduces spray coverage, while the vegetative stage offers more extensive leaf area, enhancing spray interception. However, dense canopies at later stages may impede spray penetration. Plant physiology also varies, affecting the absorption and translocation of sprayed substances (Fishel and Ferrell, 2010).

Other environmental factors come into play as well. Precipitation Pr affects the population size, with heavy precipitation leading to a decrease in the population (Ge et al., 2003). It also increases soil water content *SW* which affects the emergence rate of *H. armigera* similar to air relative humidity *RHa* (Fajun et al., 2003). The presence of fruiting organs during the plant growth stage *PGS* is important for population dynamics, as it serves as the oviposition site for females (Fitt, 1989). Crop variety *V*, such as transgenic Bt cotton, can suppress the second generation of *H. armigera*, while both different cropping systems *CS* and adjacent crops *AC* can influence the population structure (Wardhaugh et al., 1980; Gao et al., 2010; Lu et al., 2013). Finally, wind *W* and wind direction play a significant role in the emergence of *H. armigera*, influencing the distance covered during migration from nearby locations. Additionally, wind conditions at the time of spraying W_s can also impact the effectiveness of the intervention. These various factors collectively shape the population dynamics of *H. armigera* in a complex and interconnected manner as defined through domain knowledge and depicted in the causal graph (Figure 3).

4.2. Invariant & causal learning for robust pest prediction

Our goal is to predict near-future pest populations (Y_{t+1}) using Earth observation (EO) and environmental data (X_t) along with weather forecasts (W_{t+1}) by learning the function $y_{t+1} = f(x_t, w_{t+1})$. Pest management recommendations heavily depend on these predictions. Conventional machine learning methods (Aparecido et al., 2019; Skawsang et al., 2019; Xiao et al., 2019; Zhang et al., 2019), which often assume that data points are independent and identically distributed (i.i.d.), struggle to generalize to unseen environments, capture spatiotemporal variability, and adapt to climate change. These methods are prone to learning spurious correlations, limiting their effectiveness in dynamic and non-i.i.d. scenarios.

To address these challenges, we turn to causal learning (Schölkopf and von Kügelgen, 2022), which leverages domain knowledge and is grounded in the principle of independent causal mechanisms. This



Figure 4. Invariant learning for robust predictions. Stable and accurate predictions in diverse environments, such as when H. armigera feeds on different crops exhibiting variations in phenotype, agricultural management practices, and spatial distribution. Traditional ML methods risk capturing spurious correlations, such as associating pest abundance with a specific crop (e.g., cotton) due to its higher frequency in the dataset, leading to biased predictions based on the underlying crop rather than true pest presence.

principle suggests that joint probabilities can be decomposed into separate mechanisms, each reflecting an underlying causal relationship that remains stable despite environmental changes. By incorporating this principle, our models can improve generalization and robustness across varying conditions.

We achieve this by integrating invariant learning with causality and categorizing dataset units into environments *E* as different agroclimatic zones or host crops (Figure 4). While *E* influences feature x_t, w_{t+1} , it does not directly affect the target Y_t . Utilizing Invariant Causal Prediction (ICP) (Heinze-Deml et al., 2018), Directed Acyclic Graphs (DAGs), and Invariant Risk Minimization (IRM) (Arjovsky et al., 2019), we can select causal features, identify potential causal relationships, and capture latent causal structures. These tools allow us to build models that are effective in current conditions and adaptable to future environmental changes.

4.3. Explainability & counterfactual reasoning for short-term advice

We define the problem as a binary classification of pest presence or absence at the next time step, using Earth observation (EO) data (X_t) and weather forecasts (W_{t+1}). The goal is to predict the pest population value at the next time step, Y_{t+1} , by learning the function $y_{t+1} = f(x_t, w_{t+1})$. To enhance the trustworthiness of our predictions, we employ Explainable Boosting Machines (EBM) (Nori et al., 2019). This glass-box model achieves high performance while providing inherent explanations at both global and local levels. EBM's additive nature allows for the sorting and visualization of feature contributions on a local scale for each one of predictions and a global level to summarize the general behavior of the model depending on features (Figure 5), which facilitates a better understanding of the primary drivers of the model and enhances trust in its outputs.

We propose generating counterfactual examples as recommended interventions to bolster trust further and provide actionable insights. Following the setup of (Mothilal et al., 2020), we search for minimal perturbations to the feature values (x_t , w_{t+1}) that would alter the prediction to the desired class using the



Figure 5. Explainability for trustworthiness enhancement, on the right, with local and global explanations of each prediction and general model behavior, respectively, & Counterfactual explanations as agricultural actionable recommendations on the left.

same model f. These counterfactual examples represent proposed actions that could be implemented in natural farm systems, ensuring practicality and feasibility (Wachter et al., 2017; Mothilal et al., 2020). The approach ensures that the generated counterfactuals are close to the original input but predicted in the desired class, providing feasible and actionable recommendations for IPM (Figure 5).

4.4. Heterogeneous treatment effects for long-term advice

We provide long-term pest prevention and suppression advice by assessing how agricultural practices (e.g., crop rotation, balanced fertilization, sowing dates) impact pest harmfulness and yield indices. Since different agro-environments may respond variably to the same practice, it is crucial to account for this heterogeneity. We estimate the conditional average treatment effect (CATE) following the potential outcomes framework (Rubin, 2005).

The CATE quantifies the difference in potential outcomes, represented as $\mathbb{E}[Y(T=1) - Y(T=0)|X]$, where Y(T) denotes the value of a random variable Y (e.g., pest harmfulness and yield) if a unit is treated with treatment $T \in \{0, 1\}$. By controlling for field characteristics X—which capture the heterogeneity across different agro-environmental conditions—we can better understand how specific practices affect outcomes in various contexts (Figure 6). This approach allows us to provide tailored and effective long-term IPM advice sensitive to each field's unique conditions (Giannarakis et al., 2022).

4.5. Causal inference for evaluating advice effectiveness

We employ causal inference techniques to assess the effectiveness of our pest control recommendations, adapting approaches recently introduced in agricultural contexts (Tsoumas et al., 2023). Specifically, in the case of pest management and with available panel data (Table 1), we utilize causal models such as difference-in-differences (DiDs) (Abadie, 2005), synthetic control (Arkhangelsky et al., 2021) and synthetic DiDs (Abadie, 2021) to quantify the treatment effect of adhering to our framework's recommendations (*treated units*) compared to those who did not (*control units*). Historical intervention data retrospectively annotated based on whether our framework recommended action, will serve as the basis for advice evaluation. Causal inference will be performed per-environment to ensure comparability between treatment and control groups, adhering to the parallel trends assumption (Lechner et al., 2011).



Figure 6. Conditional Average Treatment Effect (CATE) is seen as long-term personalized guidance. By accounting for each land unit's unique characteristics, we can estimate a distinct treatment effect for each land unit. For example, how differences in land's characteristics can change the impact of fertilizer application on increasing the risk of pest emergence in the future.

However, digital agriculture requires a two-level evaluation of interventions to disentangle the effectiveness resulting from the accuracy of the recommendation (for intervention) in terms of spacetime from the inherent efficacy of the intervention. It is crucial to determine what effect, if any, is attributable to the space and time of application and what is due to the pesticide itself.

In this context, we conducted an initial analysis using the aforementioned panel data to quantify the impact of pesticide application on pest abundance in a real-world setting without expert system guidance, employing staggered DiDs with fixed effects (Eq. 4.1).

The staggered approach accounts for units receiving treatment at different periods. We include unitfixed effects to control for each unit's time-invariant characteristics and time-fixed effects to capture overall time trends that affect all units in each period. The unit of analysis is the plot where the pest trap is located, with periods modeled at the weekly level. Here, Y_{it} represents the outcome variable, accumulated pest abundance, for each unit *i* at the time *t*, and treated_time_{it} is an indicator of whether the unit *i* receives treatment (pesticide application) in a period *t* (in a staggered manner across units). Specifically, β_0 is the intercept, β_1 is the treatment effect coefficient, α_i represents unit fixed effects, γ_t captures time fixed effects, and ε_{it} is the error term. Thus, β_1 provides the average causal effect of the treatment (pesticide application) on the outcome (accumulated pest abundance) for treated units (ATT), as presented in Table 3 for each cultivation period from 2019 to 2022.

$$Y_{it} = \beta_0 + \beta_1 \cdot \text{treated_time}_{it} + \alpha_i + \gamma_t + \epsilon_{it}$$
(4.1)

For the years 2021 and 2022, we observe a statistically significant reduction in pest abundance, while for 2019 and 2020, we find the opposite effect. At first glance, this contradiction may seem unusual, but several reasonable explanations could account for it. Since the data come from real-world agricultural practice, it likely encapsulates some of the following issues: (i) Some interventions may have been applied incorrectly regarding timing and method, reducing or eliminating their efficacy in the pest-infested plots. This could lead to a biased estimate that the pest population increased after pesticide application (Figure 7). This occurs because the counterfactual is constructed by taking the growth trend from a plot without intervention, which might not experience the same infestation or pest pressure level. So, a mistreated plot that probably follows a steeper population increase, simply due to its higher infestation levels, can lead to this fallacy that pesticide application increases pest population. (ii) After discussions with the data provider (Corteva Agriscience Hellas), noise within the control group labels may be

Staggered DiDs estimates with fixed effects							
Year	ATT	CI	<i>p</i> -value				
2019	35.6065	(30.569, 40.644)	0.000				
2020	36.9961	(29.826, 44.166)	0.000				
2021	-13.8687	(-20.803, -6.934)	0.000				
2022	-8.5789	(-13.549, -3.609)	0.001				

 Table 3. Results of staggered DiDs with controls for unobserved heterogeneity at the unit and time levels by including fixed effects

Note: It includes point estimates, 95% confidence intervals, and p-value. Numbers represent the increase/decrease of accumulated pest catchments at the trap level after the intervention.



Figure 7. A visual example of DiDS for assessing the real-world impact of pesticide application. It demonstrates how, even when the parallel trends assumption holds in both conditions, applying an intervention (i.e., spray) at a non-recommended time can lead to unexpected effects compared to applying the intervention at the recommended time.

possible. The company is confident in the labels for treated plots, as they receive this information directly from farmers. However, they cannot be as certain about the control group. Some farmers may have applied pest control practices in their plots but chose not to report them for various reasons, such as using less expensive pesticides from competitor companies or participating in eco-schemes prohibiting pesticide use. Consequently, we face a scenario of positively labeled and unlabeled data, a common issue in machine learning. (iii) The assumption of parallel trends may not hold universally, or unobserved confounders may vary over time and between units.

In a more robust causal analysis, we can technically or conceptually address these issues. Technically, we could retrospectively employ a recommendation system or consult experts, as aforementioned, to annotate each time–space slot as favorable or unfavorable for intervention. On the other hand, we can conceptually accept reality and precisely define what causal effect we retrieve. In this case, the ATT in a

real-world setting includes different application accuracy levels, farmer's skills, expert guidance, and proper timing. To address the second issue, we plan to use Positive-Unlabeled (PU) learning methods (Bekker and Davis, 2020) to train a classifier on covariates, as they are outlined in Section 3. Using the positively labeled (treated) units only as ground truth and PU learning for training, this classifier will help establish a control group consisting only of unlabeled units that are classified there with high confidence. Lastly, a formal investigation with statistical tests is required to retain only cases where the parallel trends assumption holds. Clear assumptions statements should also be made regarding the potential of unobserved confounders that may vary by time and unit. By leveraging these techniques, we aim to rigorously evaluate the impact of our recommendations on pest control outcomes and attribute the effects to the right factors, providing robust evidence for the effectiveness of our framework in diverse agricultural environments.

5. Conclusions

In conclusion, this article presents a new framework integrating causality and explainability into digital agriculture, with a focus on enhancing pest management practices. By leveraging advanced data analysis techniques, such as causal inference and invariant learning, our approach addresses the limitations of conventional correlation-based models, providing more robust and transparent decision-making tools. This framework not only supports real-time pest control interventions but also facilitates strategic long-term planning by offering insights into the heterogeneous effects of various agricultural practices.

Our study illustrates how incorporating explainability can bolster farmers' trust and adoption of sustainable practices like IPM. The framework's use of counterfactual reasoning and explainable predictions ensures that farmers receive actionable, field-specific recommendations that can adapt to different environmental conditions. Additionally, the causal analysis embedded within our methodology allows for ongoing evaluation of the framework's effectiveness, ensuring the recommendations are impactful and contribute positively to agricultural sustainability.

We consider that a successful application to pest management will highlight, in a tangible way, the broader potential of this framework to enhance digital agriculture to drive sustainable, evidence-based practices across agriculture. Therefore, we plan to implement the proposed ideas outlined in Section 4 using the data described in Section 3. In parallel, we are gathering additional in-situ data in collaboration with Corteva Agriscience Hellas to enrich our dataset for the same pest and crop, as well as independently for other crops and pests. Finally, we explore how this approach could be adapted to related areas.

Future research will aim to expand this framework beyond pest management, exploring its potential applications in other areas of digital agriculture, such as crop disease management and nutrient optimization. Additionally, integrating advanced machine learning models to account for real-time weather data and unforeseen environmental factors will further refine prediction accuracy. Developing user-friendly tools and interfaces that facilitate farmer interaction with these data-driven insights will be critical to fostering widespread adoption.

The growing demand for sustainable agriculture underlines the importance of integrating advanced data analysis frameworks like ours. By systematically quantifying and explaining agricultural interventions, this framework offers a promising pathway for enhancing the adoption of digital agriculture in alignment with global sustainability goals. This comprehensive, data-driven approach promises to make sustainable agricultural practices more practical, facilitating a transition to a resilient and environmentally conscious food system.

Open peer review. To view the open peer review materials for this article, please visit http://doi.org/10.1017/eds.2025.14.

Acknowledgments. We express our gratitude to Corteva Agriscience Hellas, particularly to Dr. George Zanakis, the Marketing & Development Manager, for their invaluable support, insights, trust, and provision of data.

Author contribution. Conceptualization: Ilias Tsoumas, Vasileios Sitokonstantinou; Methodology: Ilias Tsoumas, Vasileios Sitokonstantinou, Evagelia Lampiri; Software: Ilias Tsoumas; Formal analysis: Ilias Tsoumas; Investigation: Evagelia Lampiri;

e23-12 Ilias Tsoumas et al.

Data curation: Ilias Tsoumas; Writing—Original Draft: Ilias Tsoumas, Vasileios Sitokonstantinou, Evagelia Lampiri; Writing—Review & Editing: all authors; Visualization: Ilias Tsoumas All authors approved the final submitted draft.

Competing interests. The authors declare none.

Data availability statement. Data availability is governed by the terms of the Memorandum of Understanding (MoU) between the National Observatory of Athens and Corteva Agriscience Hellas. Access to the data can be granted upon a formal written request to both entities. Corteva Agriscience Hellas reserves the right to make the final decision regarding access to the raw in-situ data.

Ethics statement. The research meets all ethical guidelines, including adherence to the legal requirements of the study country.

Funding statement. This research is funded by the European Union's Horizon Europe Programme under Grant Agreement No. 101135422 (UNIVERSWATER—Ilias Tsoumas, Charalampos Kontoes) and supported by the European Union Horizon Europe Research and Innovation program under Grant Agreement #101070496 (Smart Droplets—Ioannis N. Athanasiadis). Vasileios Sitokonstantinou and Gustau Camps-Valls acknowledge support from the Generalitat Valenciana and the Conselleria d'Innovació, Universitats, Ciència i Societat Digital, through the project "AI4CS: Artificial Intelligence for Complex Systems: Brain, Earth, Climate, Society" (CIPROM/2021/56). Additionally, Gustau Camps-Valls acknowledges the support of the European Research Council (ERC) under the ERC Synergy Grant USMILE (Grant Agreement 855187).

References

Abadie A (2005) Semiparametric difference-in-differences estimators. The Review of Economic Studies 72(1), 1–19.

- Abadie A (2021). Using synthetic controls: Feasibility, data requirements, and methodological aspects. *Journal of Economic Literature* 59(2), 391–425.
- Aparecido LE, Rolim G, Moraes J, Costa C and Souza P (2019) Machine learning algorithms for forecasting the incidence of coffee arabica pests and diseases. *International Journal of Biometeorology* 64(4), 671–688. https://doi.org/10.1007/s00484-019-01856-1
- Arjovsky M, Bottou L, Gulrajani I and Lopez-Paz D (2019) Invariant risk minimization. arXiv preprint arXiv:1907.02893.
- Arkhangelsky D, Athey S, Hirshberg DA, Imbens GW and Wager S (2021) Synthetic difference-in-differences. American Economic Review 111(12), 4088–4118.
- Audsley E, Stacey K, Parsons DJ and Williams AG (2009) Estimation of the Greenhouse Gas Emissions from Agricultural Pesticide Manufacture and Use. Cranfield University
- Balasundram SK, Shamshiri RR, Sridhara S and Rizan N (2023) The role of digital agriculture in mitigating climate change and ensuring food security: An overview. *Sustainability* 15(6), 5325.
- Ballabio C, Panagos P and Monatanarella L (2016) Mapping topsoil physical properties at European scale using the Lucas database. *Geoderma 261*, 110–123.
- Barzman M, Bàrberi P, Birch ANE, Boonekamp P, Dachbrodt-Saaydeh S, Graf B, Hommel B, Jensen JE, Kiss J, Kudsk P, et al (2015) Eight principles of integrated pest management. *Agronomy for Sustainable Development* 35, 1199–1215.
- Basso B and Antle J (2020) Digital agriculture to design sustainable agricultural systems. Nature Sustainability 3(4), 254–256.
- Bekker J and Davis J (2020) Learning from positive and unlabeled data: A survey. Machine Learning 109(4), 719–760.
- **Boedeker W, Watts M, Clausing P and Marquez E** (2020) The global distribution of acute unintentional pesticide poisoning: Estimations based on a systematic review. *BMC Public Health* 20(1), 1–19.
- Ceglar A, Zampieri M, Toreti A and Dentener F (2019) Observed northward migration of agro-climate zones in europe will further accelerate under climate change. *Earth's Future* 7(9), 1088–1101.
- de Brogniez D, Ballabio C, Stevens A, Jones R, Montanarella L and vanWesemael B (2015) A map of the topsoil organic carbon content of europe generated by a generalized additive model. *European Journal of Soil Science 66*(1), 121–134.
- Fajun C, Baoping Z and Xiaoxi Z (2003) Effects of soil mositure during pupual stage on population development of cotton bollowrm, helicoverpa armigera (hubner). Acta Ecologica Sinica 23(1), 112–121.
- Fishel FM and Ferrell JA (2010) Managing pesticide drift: Pi232/pi232, 9/2010. EDIS 2010(7).
- Fitt GP (1989) The ecology of heliothis species in relation to agroecosystems. Annual Review of Entomology 34(1), 17–53.
- Gao Y, Zhai B, et al (2010) Active temperature selection of flying helicoverpa armigera (lepidoptera: Noctuidae) moths. Acta Entomologica Sinica 53(5), 540–548.
- Ge F, Liu X, Ding Y, Wang X and Zhao Y (2003) Life-table of helicoverpa armigera in northern china and characters of population development in southern and northern china. *Ying Yong Sheng tai xue bao= The Journal of Applied Ecology 14*(2), 241–245.
- Giannarakis G, Sitokonstantinou V, Lorilla RS and Kontoes C (2022) Towards assessing agricultural land suitability with causal machine learning. In *Proceedings of the IEEE/CVF Conference on Computer Vision and Pattern Recognition*, New Orleans, LA, USA pp. 1441–1451 doi: 10.1109/CVPRW56347.2022.00150.
- Heimpel GE, Yang Y, Hill JD and Ragsdale DW (2013) Environmental consequences of invasive species: Greenhouse gas emissions of insecticide use and the role of biological control in reducing emissions. PLoS One 8(8), e72293.
- Heinze-Deml C, Peters J and Meinshausen N (2018) Invariant causal prediction for nonlinear models. Journal of Causal Inference 6(2), 20170016.
- Howe R (1967) Temperature effects on embryonic development in insects. Annual Review of Entomology 12(1), 15-42.

- Iost Filho FH, de Bastos Pazini J, Alves TM, Koch RL and Yamamoto PT (2022). How does the digital transformation of agriculture affect the implementation of integrated pest management?. *Frontiers in Sustainable Food Systems 6* https://www. frontiersin.org/journals/sustainable-food-systems/articles/10.3389/fsufs.2022.972213 doi 10.3389/fsufs.2022.972213 ISSN 2571-581X
- Kalyebi A, Sithanantham S, OverholtW, Hassan S and Mueke J (2005). Parasitism, longevity and progeny production of six indigenous kenyan trichogrammatid egg parasitoids (hymenoptera: Trichogrammatidae) at different temperature and relative humidity regimes. *Biocontrol Science and Technology 15*(3), 255–270.
- Lechner M, et al (2011) The estimation of causal effects by difference-in-difference methods. Foundations and Trends® in Econometrics 4(3), 165–224.
- Lu Z-Z, Zalucki MP, Perkins LE, Wang D-Y and Wu L-L (2013) Towards a resistance management strategy for helicoverpa armigera in bt-cotton in northwestern china: An assessment of potential refuge crops. *Journal of Pest Science* 86, 695–703.
- Maelzer D and Zalucki M (1999) Analysis of long-term light-trap data for helicoverpa spp.(lepidoptera: Noctuidae) in australia: The effect of climate and crop host plants. *Bulletin of Entomological Research 89*(5), 455–463.
- Maelzer D and Zalucki M (2000) Long range forecasts of the numbers of helicoverpa punctigera and h. armigera (lepidoptera: Noctuidae) in australia using the southern oscillation index and the sea surface temperature. *Bulletin of Entomological Research* 90(2), 133–146.
- Marty M, Spurlock F and Barry T (2010) Volatile organic compounds from pesticide application and contribution to tropospheric ozone. In Hayes' Handbook of Pesticide Toxicology. Elsevier, pp. 571–585.
- Mironidis G and Savopoulou-Soultani M (2014) Development, survivorship, and reproduction of helicoverpa armigera (lepidoptera: Noctuidae) under constant and alternating temperatures. *Environmental Entomology* 37(1), 16–28.
- Mothilal RK, Sharma A and Tan C (2020) Explaining machine learning classifiers through diverse counterfactual explanations. In Proceedings of the 2020 Conference on Fairness, Accountability, and Transparency. Association for Computing Machinery, New York, NY, USA, pp. 607–617 https://doi.org/10.1145/3351095.3372850.
- Nanushi O, Sitokonstantinou V, Tsoumas I and Kontoes C (2022) Pest presence prediction using interpretable machine learning. In 2022 IEEE 14th Image, Video, and Multidimensional Signal Processing Workshop (IVMSP), Nafplio, Greece pp. 1–5, doi: 10.1109/IVMSP54334.2022.9816284.
- Noor-ul-Ane M, Arif MJ, Gogi MD and Khan MA (2015) Evaluation of different integrated pest management modules to control helicoverpa for adaptation to climate change. *International Journal of Agriculture and Biology* 17(3).
- Nori H, Jenkins S, Koch P and Caruana R (2019) Interpretml: A unified framework for machine learning interpretability. arXiv preprint arXiv:1909.09223.
- Pearl J (2009) Causality. Cambridge University Press.
- Pereira FP, Reigada C, Diniz AJF and Parra JRP (2019). Potential of two trichogrammatidae species for helicoverpa armigera control. *Neotropical Entomology* 48, 966–973.
- Rubin DB (2005) Causal inference using potential outcomes: Design, modeling, decisions. *Journal of the American Statistical* Association 100(469), 322–331.
- Schölkopf B and von Kügelgen J (2022) From statistical to causal learning. In Proceedings of the International Congress of Mathematicians 1.
- Sharma A, Kumar V, Shahzad B, Tanveer M, Sidhu GPS, Handa N, Kohli SK, Yadav P, Bali AS, Parihar RD, et al (2019) Worldwide pesticide usage and its impacts on ecosystem. SN Applied Sciences 1, 1–16.
- Sharma PK,Kumar U,Vyas S, Sharma S,&Shrivastava S (2012) Monitoring of helicoverpa armigera (hubner)(lepidoptera: Noctuidae) through pheromone traps in chickpea (cicer arietinum) crop and influence of some abiotic factors on insect population. Journal of Environmental Science, Toxicology and Food Technology 1(5), 44–46.
- Sharma HC and Prabhakar CS (2014) Impact of climate change on pest management and food security. In Integrated Pest Management. Elsevier, pp. 23–36.
- Sharma A, Reeves M and Washburn C (2022) Pesticides and Climate Change: A Vicious Cycle (tech. rep.). Pesticide Action Network North America.
- Sitokonstantinou V, Porras EDS, Bautista JC, Piles M, Athanasiadis I, Kerner H, Martini G, Sweet L-b, Tsoumas I, Zscheischler J, et al (2024) Causal machine learning for sustainable agroecosystemsar. Xiv preprint arXiv:2408.13155.
- Skamarock WC, Klemp JB, Dudhia J, Gill DO, Liu Z, Berner J,Wang W, Powers JG, Duda MG, Barker DM, et al (2019) A Description of the Advanced Research WRF Model Version 4, vol. 145(145). Boulder, CO: National Center for Atmospheric Research, p. 550.
- Skawsang S, Nagai M, Tripathi N and Soni P (2019) Predicting rice pest population occurrence with satellite-derived crop phenology, ground meteorological observation, and machine learning: A case study for the central plain of thailand. *Applied Sciences 9*, 4846. https://doi.org/10.3390/app9224846
- Skendžić S, Zovko M, Živković IP, Lešić V and Lemić D (2021) The impact of climate change on agricultural insect pests. *Insects* 12(5), 440.
- Spokas K and Wang D (2003) Stimulation of nitrous oxide production resulted from soil fumigation with chloropicrin. Atmospheric Environment 37(25), 3501–3507.
- Tsoumas I, Giannarakis G, Sitokonstantinou V, Koukos A, Loka D, Bartsotas N, Kontoes C and Athanasiadis I (2023) Evaluating digital agriculture recommendations with causal inference. *Proceedings of the AAAI Conference on Artificial Intelligence* 37(12), 14514–14522.

- Wachter S, Mittelstadt B and Russell C (2017) Counterfactual explanations without opening the black box: Automated decisions and the GDPR. *Harvard Journal of Law & Technology 31*, 841.
- Wardhaugh K, Room P and Greenup L (1980) The incidence of heliothis armigera (hübner) and h. punctigera wallengren (lepidoptera: Noctuidae) on cotton and other host-plants in the namoi valley of new south wales. *Bulletin of Entomological Research* 70(1), 113–131.
- Xiao Q, Li W, Kai Y, Chen P, Zhang J and Wang B (2019) Occurrence prediction of pests and diseases in cotton on the basis of weather factors by long short term memory network. *BMC Bioinformatics 20*(Suppl 25), 688. https://doi.org/10.1186/s12859-019-3262-y
- Xu S, Sheng C and Tian C (2020) Changing soil carbon: Influencing factors, sequestration strategy and research direction. Carbon Balance and Management 15, 1–9.
- Zhang J, Huang Y, Pu R, González-Moreno P, Yuan L, Wu K and Huang W (2019) Monitoring plant diseases and pests through remote sensing technology: A review. *Computers and Electronics in Agriculture 165*, 104943. https://doi.org/10.1016/j. compag.2019.104943

Cite this article: Tsoumas I, Sitokonstantinou V, Giannarakis G, Lampiri E, Athanassiou C, Camps-Valls G, Kontoes C and Athanasiadis IN (2025). Leveraging causality and explainability in digital agriculture. *Environmental Data Science*, 4: e23. doi:10.1017/eds.2025.14